



A Novel Enriched Galerkin Method for Flow and Transport in Fractured Porous Media

Kadeethum, Teeratorn; Nick, Hamid M.; Lee, Sanghyun; Ballarin, Francesco; Richardson, C N.; Salimzadeh, Saeed

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kadeethum, T., Nick, H. M., Lee, S., Ballarin, F., Richardson, C. N., & Salimzadeh, S. (2019). *A Novel Enriched Galerkin Method for Flow and Transport in Fractured Porous Media*. Abstract from 2019 SIAM Conference on Mathematical & Computational Issues in the Geosciences, Houston, Texas, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

2019 SIAM Conference on Mathematical & Computational Issues in the Geosciences

Part of [MS40 Verification Benchmarks for Single-phase Flow in Three-dimensional Fractured Porous Media - Part I of II](#)
A Novel Enriched Galerkin Method for Flow and Transport in Fractured Porous Media

Abstract. Fluid flow and solute transport in fractured porous media is the backbone of many applications including groundwater flow, underground energy harvesting, earthquake prediction, and biomedical engineering. The traditional continuous Galerkin (CG) method is not suitable for the transport equation due to lack of mass conservation. The discontinuous Galerkin (DG) method mitigates this problem; however, its computational cost is considerably more than the CG method. In this study, a robust and efficient discretization method based on the interior penalty enriched Galerkin (EG) method is proposed. This method requires fewer degrees of freedom than those of the DG method, while it achieves the same accuracy. The flow and transport models of rock matrix and fractures domains are investigated in both equidimensional and mixed dimensional settings. The CG and EG methods are compared in four geometries to evaluate the accuracy and efficacy of the proposed method [Inga et al., Verification benchmarks for single-phase flow in three-dimensional fractured porous media, 2018]. The results illustrate the superiority of the EG method in solving the flow and transport equations in fractured porous media. Furthermore, the computational burden of the EG method is not significant.

Authors

- Teeratrorn Kadeethum, Technical University of Denmark, Denmark, teekad@dtu.dk
- Hamid M. Nick, Technical University of Denmark, Denmark, hamid@dtu.dk
- Sanghyun Lee, Florida State University, U.S., lee@math.fsu.edu
- Francesco Ballarin, SISSA-ISAS International School for Advanced Studies, Italy, francesco.ballarin@sissa.it
- C N. Richardson, University of Cambridge, United Kingdom, chris@bpi.cam.ac.uk
- Saeed Salimzadeh, Technical University of Denmark, Denmark, saeeds@dtu.dk

[GS19 Home](#)[Program](#)[Speaker Index](#)[Hotel](#)[Transportation](#)[Registration](#)

SIAM Conference Participation System

Corrections or problems using this system? Email meetings@siam.org.

Bug reports to duggan@siam.org.